

SUGGESTED TIMES OF ADMINISTRATION

acarbose	Precose	7:30am	AC/WM	metformin/glipizide	Metaglip	7:30am	WM
Achromycin	tetracycline	6:30am	AC	metformin/glyburide	Glucovance	7:30am	WM
alendronate	Fosamax	6:30am	AC	metoclopramide	Reglan	6:30am	AC
amoxicillin/clauvulanic	Augmentin	9.1.5	WM	miglitol	Glyset	7:30am	AC/WM
Axid, Pepcid, Tagamet, Zantac	H2 antagonists	5/9pm	PM	mycophenolate	Cellcept	6:30a-4:30p	AC
calcium acetate	PhosLo	9.1.5	WM	netaglinide	Starlix	7:30am	AC/WM
captopril	Capoten	6:30am	AC*	nifedipine LA	Procardia-XL	6:30am	AC*
cefuroxime	Ceftin	9.5	WM	nitrofurantoin	Macrobid	9.1.5	WM
Celebrex, Motrin, Relafen	NSAIDS	5.1.9.	WM	no more than 15" AC	Novolog/Humalog"log"	7:00am	AC/WM
chlorpropamide	Diabinese	7:30am	WM	no more than 30" AC	Regular insulin "lin"	6:30am	AC
cilostazol	Pletal	6:30am	AC	no more than 60" AC	NPH insulin "lin"	6:30am	AC
ciprofloxacin	Cipro	6:30am	AC	omeprazole	Prilosec	6:30am	AC
esomeprazole	Nexium	6:30am	AC	Omnipan	ampicillin	6:30am	AC
famotidine	Pepcid	5/9pm	PM	pantoprazole	Protonix	9:00am	ANY
glimepiride	Amaryl	7:30am	WM	PenVK	penicillin	6:30am	AC
glipizide	Glucotrol	6:30am	AC	pioglitazone	Actos	9:00am	ANY
glyburide	Diabeta	7:30am	WM	rabeprazole	Aciphex	9:00am	ANY
glyburide	Micronase	7:30am	WM	ranitidine	Zantac	5/9pm	PM
glyburide micronized	Glynase	7:30am	WM	repaglinide	Prandin	7:30am	AC/WM
ibandronate	Boniva	6:30am	AC	risedronate	Actonel	6:30am	AC
insulin detemir	Levimir	9:00am	ANY	rosiglitazone	Avandia	9:00am	ANY
insulin glargine	Lantus	9:00am	ANY	selgiline	Eldepryl	9.1	ANY
lansoprazole	Prevacid	6:30am	AC	sevelamer carbonate	Renvela	9.1.5	WM
lanthanum carbonate	Fosrenol	9.1.5	WM	sevelamer HCl	Renagel	9.1.5	WM
levofloxacin	Levaquin	6:30am	AC	sitagliptin	Januvia	9:00am	ANY
l-thyroxine	Synthroid	6:30am	AC*	sitagliptin/metformin	Janumet	9:00am	ANY
may give with largest meal	"statins"	5/9:00pm	PM	sulcrafate	Carafate	6:30am	AC
metformin	Fortamet	9:00am	ANY	tolazemide	Tolinase	6:30am	AC
metformin	Glucophage	7:30am	WM	tolbutamide	Orinase	6:30am	AC
metformin	Riomet	7:30am	WM	warfarin	Coumadin	5/9:00pm	PM

“AC” - on an empty stomach, 1 hour before or 2 hours after a meal
 * - “AC” not required, dose can be adjusted if given w/food check with resident as to what time they have been taking this medication prior to admission, may be given at 9:00am

QD - once a day, any time of day is acceptable according to resident preferences or physician specific orders
 PC - up to 2 hours after a meal
 WM - with or up to 2 hours after a meal

