

SUGGESTED TIMES OF ADMINISTRATION

Actonel	risedronate	6:30am	AC	Levemir	insulin detemir	9:00am	ANY
Aciphex	rabeprazole	9:00am	ANY	Macrobid	nitrofurantoin	9.1.5	WM
Actos	pioglitazone	9:00am	ANY	Metaglip	metformin/glipizide	7:30am	WM
Amaryl	glimepiride	7:30am	WM	Micronase	glyburide	7:30am	WM
ampicillin	Omnipran	6:30am	AC	Nexium	esomeprazole	6:30am	AC
Augmentin	amoxicillin/clauvulanic	9.1.5	WM	Novolog/Humalog"log"	no more than 15" AC	7:00am	AC/WM
Avandia	rosiglitazone	9:00am	ANY	NPH insulin "lin"	no more than 60" AC	6:30am	AC
Boniva	ibandronate	6:30am	AC	NSAIDS	Celebrex, Morin, Relafen, etc	5.1.9.	WM
Capoten	captopril	6:30am	AC*	Orinase	tolbutamide	6:30am	AC
Carafate	sulcrafate	6:30am	AC	penicillin	PenVK	6:30am	AC
Ceftin	cefuroxime	9.5	WM	Pepcid	famotidine	5/9pm	PM
Cellcept	mycophenolate	6:30a-4:30p	AC	PhosLo	calcium acetate	9.1.5	WM
Cipro	ciprofloxacin	6:30am	AC	Pletal	cilostazol	6:30am	AC
Coumadin	warfarin	5/9:00pm	PM	Prandin	repaglinide	7:30am	AC/WM
Diabeta	glyburide	7:30am	WM	Precose	acarbose	7:30am	AC/WM
Diabinese	chlorpropamide	7:30am	WM	Prevacid	lansoprazole	6:30am	AC
Eldepryl	selgiline	9.1	ANY	Prilosec	omeprazole	6:30am	AC
Fortamet	metformin	9:00am	ANY	Procardia-XL	nifedipine LA	6:30am	AC*
Fosamax	alendronate	6:30am	AC	Protonix	pantoprazole	9:00am	ANY
Fosrenol	lanthanum carbonate	9.1.5	WM	Reglan	metoclopramide	6:30am	AC
Glucophage	metformin	7:30am	WM	Regular insulin "lin"	no more than 30" AC	6:30am	AC
Glucotrol	glipizide	6:30am	AC	Renagel	sevelamer HCl	9.1.5	WM
Glucovance	metformin/glyburide	7:30am	WM	Renvela	sevelamer carbonate	9.1.5	WM
Glynase	glyburide micronized	7:30am	WM	Riomet	metformin	7:30am	WM
Glyset	miglitol	7:30am	AC/WM	Starlix	netaglinide	7:30am	AC/WM
H2 antagonists	Axid, Pepcid, Tagamet, Zantac	5/9pm	PM	"statins"	may give with largest meal	5/9:00pm	PM
Janumet	sitagliptin/metformin	9:00am	ANY	Synthroid	l-thyroxine	6:30am	AC*
Januvia	sitagliptin	9:00am	ANY	tetracycline	Achromycin	6:30am	AC
Lantus	insulin glargine	9:00am	ANY	Tolinase	tolazemide	6:30am	AC
Levaquin	levofloxacin	6:30am	AC	Zantac	ranitidine	5/9pm	PM

“AC” - on an empty stomach, 1 hour before or 2 hours after a meal
 * - “AC” not required, dose can be adjusted if given w/food check with resident as to what time they have been taking this medication prior to admission, may be given at 9:00am

QD - once a day, any time of day is acceptable according to resident preferences or physician specific orders

PC - up to 2 hours after a meal

WM - with or up to 2 hours after a meal

